

Application of Sumudu Transform to Fractional Integro-Differential Equations Involving Generalized R -Function

Shweta Shrivastava¹, Dinesh Singh Tomar² and Ashish Verma³

¹ *Department of Applied Science
Madhav Institute of Technology and Science
Gwalior (M.P.), India
shwetashrivastava986@gmail.com*

² *School of Engineering and Technology
ITM University, Gwalior (M.P.), India
dinesh2006_tomar@yahoo.co.in*

³ *Department of Mathematics,
Prof. Rajendra Singh (Rajju Bhaiya) Institute of Physical Sciences,
VBS Purvanchal University, Jaunpur (UP), India
vashish.lu@gmail.com*

Abstract

In this paper, we applied Sumudu transform to investigate the solution of a Volterra type fractional integro-differential equation involving generalized R -function. Some specific cases are also mentioned.

Keywords: Sumudu transform, Volterra type Integro-differential equation, Riemann-Liouville Fractional Integral and Derivative, Generalized R -function of Lorenzo-Hartley.

1 Introduction

With the arrival of computers and advanced softwares, the differential equations played an essential role in all aspects of applied mathematics and gained greater importance. Numerous integral based transforms such as Laplace, Mellin, Fourier, Hankel etc. are applied to solve differential equations dealing with the engineering problems. In 1993, Watugala [7] proposed the Sumudu transform. Unit and scale preserving properties of Sumudu transform has a marvelous potential of applicability in the field of engineering mathematics and applied sciences. Weerakoon [11] gave the inverse Sumudu transform. We can find further details about it from Belgacem et al [5, 6], Loonker and Banerji [4] and many others.

The aim of this paper is to present some of the significant characteristics of Sumudu transform and a simple alternative derivation of the solution of Volterra type fractional integro-differential equation. Many authors have demonstrated the solution of fractional-integro-differential equation, including Barrett [8], Kilbas, Saigo, and Saxena [1], Ross and Sachdeva [2], Saxena [9], Jain and Tomar [10] and others.

2 Preliminaries

2.1 Sumudu Transform:

Over the set of function

$$(2.1) \quad A = \{f(t) | \exists M, \tau_1, \tau_2 > 0, |f(t)| < Me^{\frac{|t|}{\tau_j}}, \text{ if } t \in (-1)^j \times [0, \infty)\},$$

the Sumudu transform for the function of exponential order defined [7] by

$$(2.2) \quad G(u) = S[f(t) : u] = \begin{cases} \int_0^\infty e^{-t} f(ut) dt, & 0 \leq u < \tau_2, \\ \int_0^\infty e^{-t} f(ut) dt, & -\tau_1 \leq u < 0. \end{cases}$$

The Sumudu transform, in other words can be written as

$$(2.3) \quad G(u) = S[f(t) : u] = \frac{1}{u} \int_0^\infty e^{-\frac{t}{u}} f(t) dt, \quad u \in (-\tau_1, \tau_2).$$

The convolution of Sumudu Transform is

$$(2.4) \quad S(f * g)(\tau) = sS[f(\tau)]S[g(\tau)] = sF(s)G(s), \text{ for } \Re(s) > 0.$$

The Sumudu transform inversion formula is given by

$$(2.5) \quad f(t) = \frac{1}{2\pi i} \int_{a-i\infty}^{a+i\infty} e^{\frac{t}{u}} G(u) du.$$

2.2 Generalized R-function:

The generalized R-function is defined by, Lorenzo-Hartley [3] as:

$$(2.6) \quad G_{q,\nu,\gamma}(a, t) = \sum_{j=0}^{\infty} \frac{(\gamma)_j (a)^j t^{(\gamma+j)q-\nu-1}}{\Gamma(j+1)\Gamma(\gamma+j)q-\nu}, \quad \Re(q\gamma - \nu) > 0,$$

Which is a particular case of $\gamma = 1$, reduces it to a well-known R-function.

2.3 Fractional Integral and Derivative of Riemann-Liouville

The α order fractional integral of Riemann-Liouville is defined as

$$(2.7) \quad {}_0D_\tau^{-\alpha} h(\tau) = \frac{1}{\Gamma(\alpha)} \int_0^\tau (\tau - t)^{\alpha-1} f(t) dt, \text{ where } \Re(\alpha) > 0.$$

Similarly, the η order fractional derivative of Riemann-Liouville is defined as

$$(2.8) \quad {}_0D_\tau^\eta [h(\tau)] = \frac{1}{\Gamma(n-\eta)} \frac{d^n}{d\tau^n} \int_0^\tau (\tau - t)^{n-\eta-1} f(t) dt, \text{ where } \Re(\eta) > 0.$$

The Sumudu transform of fractional derivative is defined as [2, 11]

$$(2.9) \quad S[{}_0D_\tau^\eta (h(\tau); s)] = s^{-\eta} H(s) - \left[\sum_{r=1}^n s^{-r} {}_0D_\tau^{\eta-r} h(\tau) \right]_{\tau=0}.$$

2.4 Sumudu Transform of Generalized R -function

Sumudu Transform of Generalized R -function is given by

$$(2.10) \quad S[G_{q,\nu,\gamma}(a, t)] = \int_0^\infty e^{-t} G_{q,\nu,\gamma}(a, st) dt,$$

$$(2.11) \quad S[G_{q,\nu,\gamma}(a, t)] = s^{\gamma q - \nu - 1} [1 - (as^q)]^{-\gamma}.$$

3 Solution of Fractional Integro-Differential Equation

Different physical phenomenon such as diffusion can be modeled in terms of integro-differential equation. These can be elongated to fractional integral equation by replacing their fractional counterparts with ordinary integral. Considering integro-differential equation of Volterra type involving generalized R -function and can be solved by using Sumudu transform.

Theorem 3.1. *Let $q, \nu, \gamma, k, \eta \in \mathbb{C}$, $0 \leq \tau \leq 1$ and $\Re(q) > 0$, $\Re(\nu) > 0$, $\Re(q - \nu) > 0$. Then the Cauchy problem for the Volterra type fractional integro-differential equation*

$$(3.1) \quad {}_0D_\tau^\alpha[h(\tau)] = \eta f(\tau) + k \int_0^\tau G_{q,\nu,\gamma}(a, t) h(\tau - t) dt$$

along with the initial conditions ${}_0D_\tau^{\alpha-k}[h(\tau)] = a_k$, $k = 1, \dots, n = -[-\Re(\eta)]$, $-1 < \eta \leq n$, $n \in \mathbb{N}$. Where a_1, \dots, a_k are prescribed constants and unique solution to the Cauchy problem (3.1) provided by

$$(3.2) \quad h(\tau) = \sum_{k=1}^n a_k \Omega_k(\tau) + \eta \int_0^\tau \theta(\tau - t) f(t) dt$$

where

$$(3.3) \quad \Omega_k(\tau) = \sum_{m=0}^{\infty} k^m G_{q,k+(\nu-\alpha)m-\alpha-1,\gamma m}(a, \tau)$$

and

$$(3.4) \quad \theta(\tau - t) = \sum_{m=0}^{\infty} k^m G_{q,(\nu-\alpha)m-\alpha-1,\gamma m}(a, \tau - t).$$

Proof. Applying Sumudu transform on equation (3.1), we have

$$(3.5) \quad s^{-\alpha} H(s) - \sum_{k=1}^n s^{-k} D_\tau^{\alpha-k} h(\tau)|_{\tau=0} = \eta F(s) + k s H(s) \frac{s^{\gamma q - \nu - 1}}{[1 - as^q]^\gamma}$$

$$H(s) = \sum_{k=1}^n s^{\alpha-k} a_k \sum_{m=0}^{\infty} k^m \frac{s^{(\gamma q - \nu + \alpha)m}}{[1 - as^q]^{\gamma m}} + \eta s^\alpha F(s) \sum_{m=0}^{\infty} k^m \frac{s^{(\gamma q - \nu + \alpha)m}}{[1 - as^q]^{\gamma m}}$$

Inverting the Sumudu transform, we have

$$(3.6) \quad s^{-1}[H(s)] = \sum_{k=1}^n a_k s^{-1} \left[\sum_{m=0}^{\infty} k^m \frac{s^{(\gamma q - \nu + \alpha)m + \alpha - k + 1 - 1}}{[1 - as^q]^{\gamma m}} \right] + \eta s^{-1} \left[F(s) \sum_{m=0}^{\infty} k^m \frac{s^{(\gamma q - \nu + \alpha)m + \alpha + 1 - 1}}{[1 - as^q]^{\gamma m}} \right]$$

Now applying the Sumudu transform theorem of Convolution

$$(3.7) \quad h(\tau) = \sum_{k=1}^n a_k \Omega_k(\tau) + \eta \int_0^{\tau} \theta(\tau - t) f(t) dt$$

where $\Omega_k(\tau)$ and $\theta(\tau - t)$ is given respectively by equation (3.3) and (3.4). □

Theorem 3.2. Let $q, \nu, \gamma, k, \eta \in \mathbb{C}$, $0 \leq \tau \leq 1$ and $\Re(q) > 0$, $\Re(\nu) > 0$, $\Re(q - \nu) > 0$. Then the Volterra type integral equation

$$(3.8) \quad {}_0D_{\tau}^{-\alpha}[h(\tau)] = \eta f(\tau) + k \int_0^{\tau} G_{q, \nu, \gamma}(a, t) h(\tau - t) dt$$

has a solution $h(\tau) = \eta \int_0^{\tau} \theta(\tau - t) f(t) dt$ where

$$(3.9) \quad \theta(\tau - t) = \sum_{m=0}^{\infty} k^m G_{q, (\nu + \alpha)m + \alpha - 1, \gamma m}(a, \tau - t).$$

Proof. Applying Sumudu transform on equation (3.8), we have

$$(3.10) \quad \begin{aligned} s^{\alpha} H(s) &= ksH(s) \frac{s^{\gamma q - \nu - 1}}{[1 - as^q]^{\gamma}} + \eta F(s) \\ H(s) &= \eta s^{-\alpha} F(s) \sum_{m=0}^{\infty} k^m \frac{s^{(\gamma q - \nu - \alpha)m}}{[1 - as^q]^{\gamma m}} \end{aligned}$$

Inverting the Sumudu transform, we have

$$(3.11) \quad s^{-1}[H(s)] = s^{-1} \left[\eta s^{-\alpha} F(s) \sum_{m=0}^{\infty} k^m \frac{s^{(\gamma q - \nu - \alpha)m}}{[1 - as^q]^{\gamma m}} \right]$$

Now applying the Sumudu transform theorem of Convolution, we get

$$(3.12) \quad h(\tau) = \eta \int_0^{\tau} \theta(\tau - t) f(t) dt,$$

where $\theta(\tau - t)$ is given by (3.11). □

4 Conclusion:

In the present paper we have utilized an interesting and efficient transform to obtain solution of a fractional integro-differential equation. The results derived in this paper are expected to apply to a wide range of areas including mathematical, engineering, physical and chemical sciences.

References

- [1] A. A. Kilbas, M. Saigo and R. K. Saxena, Solution of Volterra integro-differential equations with generalized Mittag-Leffler functions in the kernels, *J. Integral Equ. and Appl.*, 14 (2002), 377-396.
- [2] B. Ross and B. Sachdeva, The solution of certain integral equations by means of operators of arbitrary order, *Amer. Math. Monthly* 97 (1990), 498-503.
- [3] C.F. Lorenzo and T.T. Hartley, Generalized functions for the fractional calculus , *NASA, Tech. Pub.* 209424 (1999), 1-17.
- [4] D. Loonker and P.K. Banerji, On the solution of distributional Abel integral equation by distributional Sumudu transform, *Int. J. of Maths and Mathematical Sciences*, 2011, 1-8.
- [5] F. B. M. Belgacem and A. A. Karaballi, and S.L. Kalla, Analytical investigations of Sumudu transform and applications to integral equations, *Mathematical Prob. In Engg.*, 2003, 103-118.
- [6] F. B. M. Belgacem and A. A. Karaballi, Sumudu transform fundamental properties investigations and applications, *International Journal of Appl. Maths. and Stochastic Anal.*, 2006, 1-23.
- [7] G. K. Watugala, Sumudu transform: a new integral transform to solve differential equations in control engineering problems, *Int. J. Math. Educ. Sci. Technol.* 24 (1993), 35-43.
- [8] J. H. Barrett, Differential equations of non integer order, *Canad. J. Math.*, 6 (1954) 429-541.
- [9] R. K., Saxena, Alternative derivation of the solution of certain integro- differential equations of Volterra type, *Ganita Sandesh*, 17 (2003), 51-56.
- [10] Renu Jain and Dinesh Singh Tomar, An Integro-Differential Equation Of Volterra Type With Sumudu Transform, *Mathematica Aeterna* , 2 (2012), 541-547.
- [11] S. Weerakoon, Application of Sumudu transform to partial differential equations, *Int. J. of Mathematical Edu. in Sci. and Technology* 25 (1994) 277-283.